



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

we have, by multiplying all such inequalities from $n = 2$ to $n = n$,

$$b_n > \frac{1}{n}$$

and therefore the b series is divergent and hence also the a series.

Also solved by E. H. CLARKE, R. A. JOHNSON, H. L. OLSON, ARTHUR PELLETIER, and the Proposer.

2758 [1919, 124]. Proposed by LEONARD RICHARDSON, University of British Columbia.
Prove that, if r be a positive integer,

$$\int_0^{\pi/2} \frac{\sin (2r+1)\psi}{\sin \psi} d\psi = \frac{\pi}{2}.$$

and

$$\int_0^{\pi/2} \frac{\sin 2r\psi}{\sin \psi} d\psi = 2 \left\{ 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots + \frac{(-1)^{r-1}}{2r-1} \right\}.$$

I. SOLUTION BY ELIJAH SWIFT, University of Vermont.

We have the identity

$$\sin n\psi - \sin (n-2)\psi = 2 \sin \psi \cos (n-1)\psi,$$

or

$$\sin n\psi = 2 \sin \psi \cos (n-1)\psi + \sin (n-2)\psi.$$

This yields us an easy reduction formula for the two integrals. For the second,

$$\begin{aligned} \int_0^{\pi/2} \frac{\sin 2r\psi}{\sin \psi} d\psi &= 2 \int_0^{\pi/2} \cos (2r-1)\psi \cdot d\psi + \int_0^{\pi/2} \frac{\sin (2r-2)\psi}{\sin \psi} d\psi \\ &= 2 \frac{(-1)^{r-1}}{2r-1} + \int_0^{\pi/2} \frac{\sin (2r-2)\psi}{\sin \psi} d\psi. \end{aligned}$$

Applying this r times, we reach the indicated result.

For the first integral,

$$\int_0^{\pi/2} \frac{\sin (2r+1)\psi}{\sin \psi} d\psi = 2 \int_0^{\pi/2} \cos 2r\psi \cdot d\psi + \int_0^{\pi/2} \frac{\sin (2r-1)\psi}{\sin \psi} d\psi.$$

The first of these is 0. Repeating this process r times, the required integral

$$= \int_0^{\pi/2} \frac{\sin \psi}{\sin \psi} \cdot d\psi = \frac{\pi}{2}.$$

Solved similarly by A. M. HARDING and C. C. YEN.

II. SOLUTION BY R. D. BOHANNAN, Ohio State University.

Let $\cos \psi + i \sin \psi = z$ and $\cos \psi - i \sin \psi = 1/z$. Then, for the first integral, we have

$$\begin{aligned} \frac{\sin (2r+1)\psi}{\sin \psi} d\psi &= \frac{z^{2r+1} - \frac{1}{z^{2r+1}}}{z - \frac{1}{z}} \frac{dz}{iz}, \\ &= \left\{ z^{2r} + z^{2r-2} + \cdots + z^2 + 1 + \frac{1}{z^2} + \cdots + \frac{1}{z^{2r}} \right\} \frac{dz}{iz}. \end{aligned}$$

The first and last terms, after integration, give

$$\frac{1}{2ri} \left(z^{2r} - \frac{1}{z^{2r}} \right) \quad \text{or,} \quad \frac{1}{r} \sin 2r\psi \Big]_0^{\pi/2},$$

which is zero; likewise, all other pairs, equidistant from the ends. The middle term gives $(\log z)/i$ or ψ , since $z = e^{i\psi}$, and with the given limits, this reduces to $\pi/2$.

Similarly, the second integral comes from

$$\left\{ z^{2r-1} + z^{2r-3} + \cdots + z + \frac{1}{z} + \frac{1}{z^3} + \cdots + \frac{1}{z^{2r-1}} \right\} \frac{dz}{iz}.$$

After integration, the first and last terms, give

$$\frac{1}{i(2r-1)} \left\{ z^{2r-1} - \frac{1}{z^{2r-1}} \right\} \quad \text{or} \quad \frac{2}{2r-1} \sin (2r-1)\psi \Big|_0^{\pi/2} = \pm \frac{2}{2r-1}.$$

The result given is obtained by starting with the central terms,

$$\left(z + \frac{1}{z} \right) \frac{dz}{iz}.$$

Also solved similarly by P. J. DA CUNHA, WILLIAM HERBERG, and H. L. OLSON.

2759 [1919, 124]. Proposed by J. L. RILEY, Stephenville, Texas.

Solve the simultaneous functional equations

$$\phi(x+y) = \phi(x) + \frac{\phi(y) \cdot \psi(x)}{1 - \phi(x)\phi(y)},$$

$$\psi(x+y) = \frac{\psi(x) \cdot \psi(y)}{1 - \phi(x)\phi(y)}.$$

I. SOLUTION BY C. F. GUMMER, Kingston, Ont.

When $y = 0$ the equations show that either $\psi(x) = 0$, or $\phi(0) = 0$ and $\psi(0) = 1$. In the former case, $\phi(x)$ is constant. In the latter, the first equation gives

$$\frac{\phi(x+y) - \phi(x)}{y} = \frac{\phi(y) - \phi(0)}{y} \cdot \frac{\psi(x)}{1 - \phi(x)\phi(y)};$$

so that, if we assume that $\phi'(0)$ exists and equals a , we deduce that

$$\phi'(x) = a\psi(x).$$

If further $\psi'(0)$ exists and equals b , the second equation gives

$$\psi'(x) = \{a\phi(x) + b\}\psi(x).$$

Eliminating $\psi(x)$ between the last two equations, and writing $X(x) = a\phi(x) + b$, we get

$$X''(x) = X(x)X'(x);$$

and on integration, since $X(0) = b$ and $X'(0) = a^2$,

$$X(x) = 2A \tan (Ax + B),$$

where $\sqrt{2a^2 - b^2} = 2A$ and $b = 2A \tan B$.

Hence

$$\phi(x) = \sqrt{2} \cdot \frac{\sin Ax}{\cos (Ax + B)}, \quad \psi(x) = \frac{\cos^2 B}{\cos^2 (Ax + B)}.$$

This solution, however, fails to satisfy the functional equations unless $A = 0$. Hence the only solutions admitting derivatives for $x = 0$ are

$$\phi(x) = c, \quad \psi(x) = 0 \quad \text{and} \quad \phi(x) = 0, \quad \psi(x) = 1.$$

It is apparent that $\phi(x) = 0$, $\psi(x) = a^x$ is a solution.

Possibly the proposer intended to square the denominator on the right of the second equation. With this change and similar work we get, assuming $\phi'(0)$ and $\psi'(0)$ to exist, *either*

$$\phi(x) = c, \quad \psi(x) = 0$$